

ALU-A118 708

ALBERTA UNIV EDMONTON DEPT OF CHEMISTRY

F/G 7/4

SIMULATION OF EDGE EFFECTS IN ELECTROANALYTICAL EXPERIMENTS BY --ETC(11)

AUG 82 B SPEISER, S PONS

N00014-82-G-0017

UNCLASSIFIED

TR-8

NL

10-1
A-1
1-708



12

OFFICE OF NAVAL RESEARCH

Contract N00014-82-G-0017

Task No. NR 359-718

TECHNICAL REPORT NO. 8

Simulation of Edge Effects in Electroanalytical Experiments.
Part I. Two Dimensional Collocation and Theory for Chrono-
amperometry.

By

Bernd Speiser and Stanley Pons

Prepared for Publication
in

Canadian Journal of Chemistry

University of Alberta
Department of Chemistry
Edmonton, Alberta, Canada
T6G 2G2

August 18, 1982

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited.

DTIC
ELECTE
AUG 30 1982
H

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS THESE CANNOT BE USED FOR REPRODUCTION
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
8	AD A118 708	
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED	
Simulation of Edge Effects in Electroanalytical Experiments. I. Two Dimensional Collocation and Theory for Chronoamperometry.	Technical Report # 8	
6. AUTHOR(s)	7. PERFORMING ORG. REPORT NUMBER	
Bernd Speiser and Stanley Pons*	N00014-82-G-0017	
8. PERFORMING ORGANIZATION NAME AND ADDRESS	9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
Department of Chemistry University of Alberta Edmonton, Alberta, Canada T6G 2G2	Task No. NR 359-718	
10. CONTROLLING OFFICE NAME AND ADDRESS	11. REPORT DATE	
Office of Naval Research Chemistry Program Arlington, Virginia 22217 - Chemistry Code 472	August 18, 1982	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. SECURITY CLASS. (of this report)	
	Unclassified	
14. DISTRIBUTION STATEMENT (of this Report)		
This document has been approved for public release and sale; its distribution unlimited.		
15. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
16. SUPPLEMENTARY NOTES		
17. KEY WORDS (Continue on separate sheet if necessary and identify by block number)		
Electrochemistry, Simulation, Polynomial, Orthogonal Collocation		
18. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
The details for accurate simulations of edge effects experienced at disc electrodes in chronoamperometric experiments is given.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE
S/N 8102-11-014-6001

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

82 8 27 056

Simulation of edge-effects in electroanalytical experiments by orthogonal collocation. Part 1. Two-dimensional collocation and theory for chronoamperometry

BERND SPEISER and STANLEY PONS

Department of Chemistry, University of Alberta, Edmonton, Alta., Canada T6G 2G2
Received November 26, 1981

BERND SPEISER and STANLEY PONS, *Can. J. Chem.* 60, 1352 (1982)

A two-dimensional model to describe edge-effects at planar circular disc electrodes has been developed. Two-dimensional orthogonal collocation is used to discretize the corresponding partial differential equations. The equations for the simulation of a chronoamperometric experiment are derived.

BERND SPEISER et STANLEY PONS, *Can. J. Chim.* 60, 1352 (1982)

On a mis au point une méthode bidimensionnelle pour décrire les effets d'arête au niveau des électrodes à disques circulaires plans. On a utilisé une discrétisation orthogonale pour les équations différentielles partielles correspondantes. On a dérivé les équations de la simulation d'une expérience de chronoamperométrie.

(Traduit par le journal)

Introduction

One common simplification in the theoretical treatment of electrode processes of planar disc electrodes is the assumption of semi-infinite linear diffusion. This results in the reduction of the three-dimensional system to a one-dimensional model, where the distance x from the electrode surface is the only coordinate to be considered. Consequently, the three-dimensional form of Fick's second law

$$[1] \quad \frac{\partial c}{\partial t} = D \left[\frac{\partial^2 c}{\partial x^2} + \frac{\partial^2 c}{\partial r^2} + \frac{\partial^2 c}{\partial z^2} \right]$$

reduces to the simpler differential equation

$$[2] \quad \frac{\partial c}{\partial t} = D \frac{\partial^2 c}{\partial x^2}$$

and the mathematical and computational efforts decrease considerably. The assumption of linear diffusion to a planar electrode might imply an infinite disc without edges. For real systems, however, this of course is not possible.

Thus, measurements at finite disc electrodes should deviate from the simple one-dimensional theory. In fact, it has been shown for different electroanalytical techniques such as chronoamperometry and chronopotentiometry (1) that the diffusion current is higher than expected from the theoretical calculations based on eq. [2], i.e. the Cottrell equation (2) for chronoamperometry. For cyclic voltammograms (3) the peak separation is larger than the theoretical values given by Nicholson and Shain (4). A number of papers have accounted for these deviations as being due to additional diffusion from directions other than perpendicular to the disc electrode (see e.g. refs. 5,

6). In a system with several coupled charge transfer and chemical reactions dramatic changes of the overall voltammogram have been attributed to edge effects (7).

Several attempts have been made to calculate the effects of edge diffusion and to simulate model systems without assuming one-dimensional diffusion (5, 6). Recently, Heinze published a two-dimensional model based on the finite difference method (8, 9), while Oldham (10) and Aoki and Osteryoung (11) developed analytical solutions for the simple reversible mechanism.

These calculations showed, in agreement with experiments, that edge diffusion rates increase with increasing time and decreasing radius of the disc electrode. The deviation from pure Cottrell behavior is claimed to be between 9 and 10% for typical conditions: diffusion coefficient = 10^{-5} cm²/s, electrode radius 0.1 cm, electrolysis time 2.65 s (5, 9–11). All methods presented thus far, however, demonstrate results for only the simplest mechanisms (e.g. reversible electron transfer case). Extensions apparently require extensive computational effort.

Orthogonal collocation has been shown to provide a powerful method to simulate responses of several electroanalytical techniques to a variety of mechanisms (12–18). Up to now all these calculations have been performed using one-dimensional diffusion models. We want to report the construction of a two-dimensional model for the diffusion to and from a planar disc electrode and the discretization of the resulting second order partial differential equations to give a set of simultaneous first order ordinary differential equations.

The comparison of the results of this new model with those already presented is the subject of a

separate report now in preparation. Extensions to other mechanisms and electroanalytical experiments will also be considered.

The two-dimensional model

Consider a planar, circular disc of conducting material imbedded into an insulator (Fig. 1). The radius of the disc is r_0 . The radius r_0 of the insulator may go to infinity. From the symmetry of the problem, it follows that we may use two coordinates x (perpendicular to the disc) and r (along the radius vector of the disc). We are required to calculate the concentration c of all species as a function of x , r and the time t , where c is given by second order partial differential equations of the type (9)

$$[3] \quad \frac{\partial c}{\partial t} = D \left[\frac{\partial^2 c}{\partial x^2} + \frac{1}{r} \frac{\partial c}{\partial r} + \frac{\partial^2 c}{\partial r^2} \right] - f(c)$$

and $f(c)$ is determined by the kinetics of the system investigated. D is the diffusion coefficient.

Let us consider an EC-mechanism (4). There the differential equations describing the concentration of species A, B, and C are given by [5]–[7].



$$[5] \quad \frac{\partial c_A}{\partial t} = D_A \left[\frac{\partial^2 c_A}{\partial x^2} + \frac{1}{r} \frac{\partial c_A}{\partial r} + \frac{\partial^2 c_A}{\partial r^2} \right]$$

$$[6] \quad \frac{\partial c_B}{\partial t} = D_B \left[\frac{\partial^2 c_B}{\partial x^2} + \frac{1}{r} \frac{\partial c_B}{\partial r} + \frac{\partial^2 c_B}{\partial r^2} \right] - k_3 c_B$$

$$[7] \quad \frac{\partial c_C}{\partial t} = D_C \left[\frac{\partial^2 c_C}{\partial x^2} + \frac{1}{r} \frac{\partial c_C}{\partial r} + \frac{\partial^2 c_C}{\partial r^2} \right] + k_3 c_B$$

For $k_1 = 0$ we deal with a simple reversible electron transfer. The electrode reaction in eq. [4] is drawn as an oxidation but the extension to reductions is immediate. For simplification we may assume that the diffusion coefficients are equal:

$$[8] \quad D_A = D_B = D_C$$

This, however, is not necessary for orthogonal collocation simulations.

Depending on the electroanalytical method we wish to simulate for the mechanism chosen, certain initial and boundary conditions are imposed upon the solution of eq. [3]. Let us consider a chronoamperometric experiment, where species A, the substrate, is present at a homogeneous concentration c_A^0 at the beginning of the experiment. The product




Fig. 1. Geometry of the two-dimensional model.

species B and C may not be present at $t = 0$ (initial conditions [9], [10]).

$$[9] \quad c_A(x, r, 0) = c_A^0$$

$$[10] \quad c_B(x, r, 0) = c_C(x, r, 0) = 0$$

At the boundary $x \rightarrow \infty$, i.e. at a distance x far from the electrode where the concentrations are not changed during the experiment:

$$[11] \quad c_A(\infty, r, t) = c_A^0$$

$$[12] \quad c_B(\infty, r, t) = c_C(\infty, r, t) = 0$$

The same condition may hold for a distance r far from the electrode radially:

$$[13] \quad c_A(x, \infty, t) = c_A^0$$

$$[14] \quad c_B(x, \infty, t) = c_C(x, \infty, t) = 0$$

At the electrode surface the Nernst equation has to be fulfilled for c_A and c_B (eq. [14a]). If we step to a potential E at least 200 mV past the formal potential $E_{f,A}$ the concentration of the substrate A at the electrode is immediately forced to zero and [14a] may be replaced by [14b]. Additional conditions at

the electrode surface are given by the flux equations [15] and [16].

$$r > 0, x = 0, 0 \leq r \leq r_0$$

$$[14a] \frac{c_A(0, r, t)}{c_A(0, r_0, t)} = \exp \left[\frac{nF}{RT} (E^0 - E) \right]$$

$$[14b] c_A(0, r, t) = 0$$

$$[15] \left(\frac{\partial c_A}{\partial x} \right)_{x=0} = - \left(\frac{\partial c_A}{\partial x} \right)_{x=r_0}$$

$$[16] \left(\frac{\partial c_A}{\partial x} \right)_{x=r_0} = 0$$

At $x = 0$, but outside the electroactive area ($r > r_0$), no species may undergo electron transfer and we have boundary condition [17].

$$r > 0, x = 0, r_0 \leq r \rightarrow \infty$$

$$[17] \left(\frac{\partial c_A}{\partial x} \right)_{x=0} = \left(\frac{\partial c_A}{\partial x} \right)_{x=r_0} = \left(\frac{\partial c_A}{\partial x} \right)_{x=\infty} = 0$$

Because of the symmetry of the problem there may be no net flux of any species across the x -axis at the center of the electrode ($r = 0$).

$$r > 0, x \geq 0, r = 0$$

$$[18] \left(\frac{\partial c_A}{\partial r} \right)_{r=0} = \left(\frac{\partial c_A}{\partial r} \right)_{r=r_0} = \left(\frac{\partial c_A}{\partial r} \right)_{r=\infty} = 0$$

Equations [4]–[18] describe the two-dimensional model system for an EC-mechanism under chronoamperometric conditions.

Since there are different boundary conditions for $x = 0$ inside ($r \leq r_0$) and outside ($r > r_0$) the electrode, it is convenient to describe the whole system by two concentration functions c_1 and c_2 which are defined for $0 \leq r \leq r_0$ and $r_0 < r$ respectively. To avoid discontinuities at $r = r_0$ we have to apply some additional "boundary" conditions for all species:

$$[19] c_i(x, r_0, t) = c_j(x, r_0, t)$$

The second derivative of c with respect to r is only defined if the first derivative with respect to r has no discontinuity at $r = r_0$ and this gives the further conditions

$$[20] \left(\frac{\partial c_1}{\partial r} \right)_{r=r_0} = \left(\frac{\partial c_2}{\partial r} \right)_{r=r_0}$$

for all species.

It has been pointed out [12] that orthogonal collocation may be applied only to dimensionless systems which have been converted from semi-infinite models to models in the range 0 to 1.

In order to meet these requirements we use the following transformation equations:

$$[21] X = x/L$$

$$[22] c^* = c/c_1^0$$

$$[23a] R = r/r_0$$

$$[23b] R' = (r - r_0)/(M - r_0)$$

$$[24] \alpha = k \cdot r_0^2/D$$

$$[25] T = Dt/r_0^2$$

where X is the dimensionless distance from the electrode, L is the distance from the electrode in the x direction defined so that at $X = L$ no changes in concentration take place during the time of the experiment, c^* is the dimensionless concentration, R and R' are the dimensionless distances in the r direction inside and outside the electrode disc respectively, M is the distance in the r direction defined in the same manner as L , and T is the dimensionless simulation time.

Application of eqs. [21]–[25] and the subdivision into inner and outer functions c_1 and c_2 of eqs. [4]–[18] yield the dimensionless model [26]–[33]

$$[26] \frac{\partial c_1}{\partial T} = \beta \frac{\partial^2 c_1}{\partial X^2} + \frac{1}{R} \frac{\partial c_1}{\partial R} + \frac{\partial^2 c_1}{\partial R^2}$$

$$[27] \frac{\partial c_1}{\partial T} = \beta \frac{\partial^2 c_1}{\partial X^2} + \frac{1}{R} \frac{\partial c_1}{\partial R} + \frac{\partial^2 c_1}{\partial R^2} - \alpha c_1$$

$$[28] \frac{\partial c_2}{\partial T} = \beta \frac{\partial^2 c_2}{\partial X^2} + \frac{1}{R} \frac{\partial c_2}{\partial R} + \frac{\partial^2 c_2}{\partial R^2} + \alpha c_2$$

where

$$[29] \beta = r_0^2/L^2$$

and

$$[30] \frac{\partial c_1}{\partial T} = \beta \frac{\partial^2 c_1}{\partial X^2} + \frac{\beta'}{R' + \sqrt{\beta'}} \frac{\partial c_1}{\partial R'} + \beta' \frac{\partial^2 c_1}{\partial R'^2}$$

$$[31] \frac{\partial c_2}{\partial T} = \beta \frac{\partial^2 c_2}{\partial X^2} + \frac{\beta'}{R' + \sqrt{\beta'}} \frac{\partial c_2}{\partial R'} + \beta' \frac{\partial^2 c_2}{\partial R'^2} - \alpha c_2$$

$$[32] \frac{\partial c_2}{\partial T} = \beta \frac{\partial^2 c_2}{\partial X^2} + \frac{\beta'}{R' + \sqrt{\beta'}} \frac{\partial c_2}{\partial R'} + \beta' \frac{\partial^2 c_2}{\partial R'^2} + \alpha c_2$$

where

$$[33] \beta' = r_0^2/(M - r_0)^2$$

with the initial conditions

$$T = 0, 0 \leq X \leq 1, 0 \leq R \leq 1, 0 \leq R' \leq 1$$

$$[34] c_1(X, R, 0) = c_1(X, R', 0) = 1$$

$$[35] c_2(X, R, 0) = c_2(X, R', 0) = c_2(X, R', 0) = c_2(X, R', 0) = 0$$

and boundary conditions

$$T > 0, X = 1, 0 \leq R \leq 1$$

$$[36] c_1(1, R, T) = 1$$

$$[37] c_2(1, R, T) = c_2(1, R', T) = 0$$

$$T > 0, X = 1, 0 \leq R' \leq 1$$

$$[38] c_1(1, R', T) = 1$$

$$[39] c_2(1, R', T) = c_2(1, R, T) = 0$$

$$T > 0, 0 \leq X \leq 1, R' = 1$$

$$[40] c_1(X, 1, T) = 1$$

$$[41] c_2(X, 1, T) = c_2(X, 1, T) = 0$$

$$T > 0, X = 0, 0 \leq R \leq 1$$

$$[42a] \frac{c_1(0, R, T)}{c_1(0, R', T)} = \exp \left[\frac{nF}{RT} (E^0 - E) \right]$$

$$[42b] c_1(0, R, T) = 0$$

$$[43] \left(\frac{\partial c_1}{\partial X} \right)_{X=0} = - \left(\frac{\partial c_1}{\partial X} \right)_{X=0}$$

$$[44] \left(\frac{\partial c_2}{\partial X} \right)_{X=0} = 0$$

or

$$[53] c(x, r, t) = \lambda_{1,1} x^{N_1+1} + \lambda_{1,2} x^{N_1+2} + \dots + \lambda_{1,N_1+1} x^{N_1+1} + \lambda_{1,N_1+2} x^{N_1+2} + \dots + \lambda_{1,N_1+1} x^{N_1+1} + \lambda_{1,N_1+2} x^{N_1+2} + \dots + \lambda_{1,N_1+1} x^{N_1+1} + \lambda_{1,N_1+2} x^{N_1+2} + \dots$$

or

$$[54] c(x, r, t) = \sum_{j=1}^{N_1+1} \sum_{i=1}^{N_2+1} \lambda_{ij} x^{j-1} r^{i-1}$$

i.e. we represent the concentration by a polynomial in x and r whose time-dependent coefficients are the λ_{ij} . The polynomial is of order $N_1 + 2$ and $N_2 + 2$ with respect to x and r , respectively. The

$$T > 0, X = 0, 0 \leq R' \leq 1$$

$$[45] \left(\frac{\partial c_1}{\partial X} \right)_{X=0} = \left(\frac{\partial c_1}{\partial X} \right)_{X=0} = \left(\frac{\partial c_2}{\partial X} \right)_{X=0} = 0$$

$$T > 0, 0 \leq X \leq 1, R = 0$$

$$[46] \left(\frac{\partial c_1}{\partial R} \right)_{R=0} = \left(\frac{\partial c_1}{\partial R} \right)_{R=0} = \left(\frac{\partial c_2}{\partial R} \right)_{R=0} = 0$$

$$T > 0, 0 \leq X \leq 1, R = 1 (R' = 0)$$

$$[47] c_1(X, 1, T) = c_1(X, 0, T)$$

$$[48] c_2(X, 1, T) = c_2(X, 0, T)$$

$$[49] c_2(X, 1, T) = c_2(X, 0, T)$$

$$[50] \left(\frac{\partial c_1}{\partial R} \right)_{R=1} = -\sqrt{\beta'} \left(\frac{\partial c_1}{\partial R'} \right)_{R'=0}$$

$$[51] \left(\frac{\partial c_2}{\partial R} \right)_{R=1} = -\sqrt{\beta'} \left(\frac{\partial c_2}{\partial R'} \right)_{R'=0}$$

$$[52] \left(\frac{\partial c_2}{\partial R} \right)_{R=1} = -\sqrt{\beta'} \left(\frac{\partial c_2}{\partial R'} \right)_{R'=0}$$

We have to calculate the inner and outer concentrations of species A, B, and C respectively as a function of X , $R(R')$, and T .

Two-dimensional collocation

Orthogonal collocation uses a mathematically suitable trial function to describe the solution of a differential equation at certain points. If one uses a set of weighted orthogonal polynomials as the trial function, the differential equation is satisfied at the zeroes of the polynomial of highest order used (19, 20) for a set of coefficients which depend on time.

To perform orthogonal collocation in two-dimensions, x and r , we use a trial function depending on x and r

points in the x and r directions, respectively

$$[55] \quad c(x_i, r_k, t) = \sum_{j=1}^{N_x+1} \sum_{l=1}^{N_r+1} \lambda_{j,l} x_j^{i-1} r_l^{k-1}$$

The concentrations at the collocation points may be the elements of a concentration matrix $\Gamma_{i,k}$

$$[56] \quad \Gamma_{i,k} = X_{i,j} \cdot A_{j,l} \cdot R_{l,k}$$

which is calculated by multiplication of a matrix $X_{i,j}$, containing the x_j^{i-1} as elements, a matrix $A_{j,l}$, containing the elements $\lambda_{j,l}$, and a matrix $R_{l,k}$, containing the elements r_l^{k-1} . The superscript T denotes a transposed matrix.

Now we can calculate the first and second derivatives of the concentration at the collocation points with respect to x and r respectively:

$$[57] \quad \Gamma_{i,k}^x = X_{i,j}^x \cdot A_{j,l} \cdot R_{l,k}$$

$$[58] \quad \Gamma_{i,k}^r = X_{i,j} \cdot A_{j,l} \cdot R_{l,k}^r$$

$$[59] \quad \Gamma_{i,k}^{xx} = X_{i,j}^{xx} \cdot A_{j,l} \cdot R_{l,k}$$

$$[60] \quad \Gamma_{i,k}^{rr} = X_{i,j} \cdot A_{j,l} \cdot R_{l,k}^{rr}$$

where the superscript $x(i)$ denotes the first and accordingly $r(i)$ characterizes the second derivative with respect to $x(r)$.

From eq. [56] it follows that

$$[61] \quad A_{j,l} = X_{i,j}^{-1} \cdot \Gamma_{i,k} \cdot (R_{l,k}^T)^{-1}$$

After inserting [61] into [57] we arrive at

$$[62] \quad \Gamma_{i,k}^x = X_{i,j}^x \cdot X_{i,j}^{-1} \cdot \Gamma_{i,k} = A_{i,j} \cdot \Gamma_{i,k}$$

where the first derivative of the concentration with respect to x is expressed as a function of the concentration itself. In analogy eqs. [61] and [58] give

$$[63] \quad \Gamma_{i,k}^r = X_{i,j}^r \cdot X_{i,j}^{-1} \cdot \Gamma_{i,k} = B_{i,j} \cdot \Gamma_{i,k}$$

From [61] and [59] we get

$$[64] \quad \Gamma_{i,k}^{xx} = \Gamma_{i,k} \cdot (B_{i,j}^T)^{-1} \cdot B_{i,j}^{xx} = \Gamma_{i,k} \cdot (B_{i,j}^T)^{-1} \cdot B_{i,j}^{xx} \\ = \Gamma_{i,k} \cdot (B_{i,j}^T \cdot B_{i,j}^{xx})^{-1}$$

or

$$[65] \quad \Gamma_{i,k}^{xx} = \Gamma_{i,k} \cdot C_{i,j}$$

Analogous from [61] and [60]

$$[66] \quad \Gamma_{i,k}^{rr} = \Gamma_{i,k} \cdot (B_{i,j}^T \cdot B_{i,j}^{rr})^{-1} = \Gamma_{i,k} \cdot D_{i,j}$$

We may calculate the matrices containing the derivatives of the concentration at the collocation points from the concentration matrix once the matrices $A_{j,l}$, $B_{i,j}$, $C_{i,j}$, and $D_{i,j}$ have been determined. Returning to summation expression we

can write

$$[67] \quad \left(\frac{\partial c}{\partial x} \right)_{i,k} = \sum_{j=1}^{N_x+1} A_{i,j} c(x_j, r_k, t)$$

$$[68] \quad \left(\frac{\partial^2 c}{\partial x^2} \right)_{i,k} = \sum_{j=1}^{N_x+1} C_{i,j} c(x_j, r_k, t)$$

$$[69] \quad \left(\frac{\partial c}{\partial r} \right)_{i,k} = \sum_{l=1}^{N_r+1} B_{i,l} c(x_i, r_l, t)$$

and

$$[70] \quad \left(\frac{\partial^2 c}{\partial r^2} \right)_{i,k} = \sum_{l=1}^{N_r+1} D_{i,l} c(x_i, r_l, t)$$

As has been pointed out by Whiting and Carr (12) for the one-dimensional case, the matrices $A_{i,j}$, $B_{i,j}$, $C_{i,j}$, and $D_{i,j}$ depend solely on the type of polynomial used as trial function and may be calculated from the zeroes of this polynomial. This has to be done only once. Using the values of the elements of these matrices we are able to express the differential quotients of c with respect to x and r at all collocation points.

For linear diffusion problems it has been shown that Legendre polynomials are most suitable (12). Pons (18) used Chebyshev polynomials for diffusion under spherical geometry. It may be noted, however, that the choice of the trial function does not have a large effect on the accuracy of the solutions for higher order approximations. In general, the best trial function is determined by the symmetry of the problem.

In this work, Legendre polynomials have been used for discretization along the x -coordinate. The collocation in the direction of the r -coordinate has been performed using different types of functions including Legendre, Chebyshev, and polynomials constructed from even powers of r , which is suggested by the symmetry of the problem around $r = 0$ (19).

Discretization of the equations for chronoamperometry

We can substitute the summation expressions [67]–[70] for the partial differential quotients in eqs. [26]–[28] and [30]–[32]. This yields expressions for the time derivative of the concentration of species A, B, and C at all the collocation points as a function of the concentrations at the collocation points. If we substitute the boundary conditions [36]–[41], [42b] (i.e. we assume a step to a potential far behind E^0) and [47]–[49], we arrive at

$$[71] \quad \frac{dc_A}{dt} \Big|_{i,k} = -\beta \left[B_{i,j} c_A(x_i, r_k, T) + \sum_{l=1}^{N_r+1} B_{i,l} c_A(x_i, r_l, T) \right] \\ + \left[E_{i,j} c_A(x_i, 0, T) + E_{i,j+1} c_A(x_i, 1, T) + \sum_{l=1}^{N_r+1} E_{i,l} c_A(x_i, r_l, T) \right]$$

$$[72] \quad \frac{dc_B}{dt} \Big|_{i,k} = -\beta \left[B_{i,j} c_B(x_i, r_k, T) + \sum_{l=1}^{N_r+1} B_{i,l} c_B(x_i, r_l, T) \right] \\ + \left[E_{i,j} c_B(x_i, 0, T) + E_{i,j+1} c_B(x_i, 1, T) + \sum_{l=1}^{N_r+1} E_{i,l} c_B(x_i, r_l, T) \right] - \alpha c_B$$

$$[73] \quad \frac{dc_C}{dt} \Big|_{i,k} = -\beta \left[B_{i,j} c_C(x_i, r_k, T) + \sum_{l=1}^{N_r+1} B_{i,l} c_C(x_i, r_l, T) \right] \\ + \left[E_{i,j} c_C(x_i, 0, T) + E_{i,j+1} c_C(x_i, 1, T) + \sum_{l=1}^{N_r+1} E_{i,l} c_C(x_i, r_l, T) \right] + \alpha c_C$$

$$[74] \quad \frac{dc_A}{dt} \Big|_{i,k} = -\beta \left[B_{i,j} c_A(x_i, r_k, T) + B_{i,j+1} c_A(x_i, r_{j+1}, T) + \sum_{l=1}^{N_r+1} B_{i,l} c_A(x_i, r_l, T) \right] \\ + \beta' \left[F_{i,j} c_A(x_i, 0, T) + F_{i,j+1} c_A(x_i, 1, T) + \sum_{l=1}^{N_r+1} F_{i,l} c_A(x_i, r_l, T) \right]$$

$$[75] \quad \frac{dc_B}{dt} \Big|_{i,k} = -\beta \left[B_{i,j} c_B(x_i, r_k, T) + \sum_{l=1}^{N_r+1} B_{i,l} c_B(x_i, r_l, T) \right] \\ + \beta' \left[F_{i,j} c_B(x_i, 0, T) + \sum_{l=1}^{N_r+1} F_{i,l} c_B(x_i, r_l, T) \right] - \alpha c_B$$

$$[76] \quad \frac{dc_C}{dt} \Big|_{i,k} = -\beta \left[B_{i,j} c_C(x_i, r_k, T) + \sum_{l=1}^{N_r+1} B_{i,l} c_C(x_i, r_l, T) \right] \\ + \beta' \left[F_{i,j} c_C(x_i, 0, T) + \sum_{l=1}^{N_r+1} F_{i,l} c_C(x_i, r_l, T) \right] + \alpha c_C$$

where

$$[77] \quad E_{i,j} = \frac{C_{i,j} + R_{i,j} D_{i,j}}{R_{i,j}}$$

and

$$[78] \quad F_{i,j} = \frac{C_{i,j}}{\sqrt{\beta' + R_{i,j}}} + D_{i,j}$$

The primed terms, except T' and β' , correspond to the outer functions. Equations [71]–[76] contain $9N_x + 2N_r + 3N_c$ unknowns, i.e. $c_A(x_i, 0, T)$, $c_A(x_i, 1, T)$, $c_B(x_i, 0, T)$, $c_B(x_i, 1, T)$, $c_C(x_i, 0, T)$, $c_C(x_i, 1, T)$, $c_A(x_i, r_l, T)$, $c_B(x_i, r_l, T)$, $c_C(x_i, r_l, T)$ for all x_i ; $c_A(0, R_k, T)$, $c_B(0, R_k, T)$, $c_C(0, R_k, T)$ for all R_k ; $c_A(0, R_k', T)$, $c_B(0, R_k', T)$, $c_C(0, R_k', T)$ for all R_k' which may be computed from the boundary conditions as is shown below.

Condition [43] yields

$$[79] \quad \sum_{j=1}^{N_x+1} A_{i,j} c_A(x_j, r_k, T) = - \sum_{j=1}^{N_x+1} A_{i,j} c_B(x_j, r_k, T)$$

Expanding the sums, introducing [36], [37], and [42b], and solving for the concentration of species B at the electrode surface ($x_i = 0$) gives:

$$[80] \quad c_h(0, R_0, T) = -\frac{1}{A_{1,2}} \left[A_{1,0,1} + \sum_{i=1}^{N+1} A_{1,i} c_h(X_i, R_0, T) + c_h(X_0, R_0, T) \right]$$

Condition [44] is changed to

$$\sum_{i=1}^{N+1} A_{1,i} c_h(X_i, R_0, T) = 0$$

which gives, after introduction of [37],

$$[81] \quad c_h(0, A_0, T) = -\frac{1}{A_{1,2}} \sum_{i=1}^{N+1} A_{1,i} c_h(X_i, R_0, T)$$

In analogy it may be derived from [45] and [40] or [41]

$$[82] \quad c_h(0, R_0, T) = -\frac{1}{A_{1,1}} \left[A_{1,0,1} + \sum_{i=1}^{N+1} A_{1,i} c_h(X_i, R_0, T) \right]$$

$$[83] \quad c_h(0, R_0, T) = -\frac{1}{A_{1,1}} \sum_{i=1}^{N+1} A_{1,i} c_h(X_i, R_0, T)$$

and

$$[84] \quad c_h(0, R_0, T) = -\frac{1}{A_{1,2}} \sum_{i=1}^{N+1} A_{1,i} c_h(X_i, R_0, T)$$

The condition [46] gives, for species A, B, and C

$$[85] \quad c_h(X_0, 0, T) = -\frac{1}{C_{1,1}} \left[C_{0,1,1} c_h(X_0, 1, T) + \sum_{i=1}^{N+1} C_{1,i} c_h(X_i, R_0, T) \right]$$

$$[86] \quad c_h(X_0, 0, T) = -\frac{1}{C_{1,1}} \left[C_{0,1,1} c_h(X_0, 1, T) + \sum_{i=1}^{N+1} C_{1,i} c_h(X_i, R_0, T) \right]$$

and

$$[87] \quad c_h(X_0, 0, T) = -\frac{1}{C_{1,1}} \left[C_{0,1,1} c_h(X_0, 1, T) + \sum_{i=1}^{N+1} C_{1,i} c_h(X_i, R_0, T) \right]$$

Condition [50] yields

$$[88] \quad \sum_{i=1}^{N+1} C_{1,i} c_h(X_i, R_0, T) = \sum_{i=1}^{N+1} \sqrt{F} C_{1,i} c_h(X_i, R_0, T)$$

which may be expanded and transformed using [40] and [47]

$$[89] \quad [C_{0,1,1} c_h(X_0, 1, T) + C_{1,0,1} c_h(X_0, 0, T)] - \sum_{i=1}^{N+1} C_{1,i} c_h(X_i, R_0, T) + \sum_{i=1}^{N+1} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) + \sqrt{F} C_{0,1,1}$$

From [85] and [89] it follows that

$$[90] \quad c_h(X_0, 1, T) = c_h(X_0, 0, T) = \frac{1}{G_1} \left[\sum_{i=1}^{N+1} H_i c_h(X_i, R_0, T) + \sum_{i=1}^{N+1} C_{1,i} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) + C_{1,0} \sqrt{F} C_{0,1,1} \right]$$

where

$$[91] \quad G_1 = [C_{0,1,1} c_h(X_0, 1, T) - \sqrt{F} C_{1,1} C_{1,0} - C_{1,0,1} c_h(X_0, 0, T)]$$

Accession For	
NTIS Grant	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability Cod	
Dist	Avail and/or
A	Special



and

$$[92] \quad H_i = C_{1,0,1} C_{1,i} - C_{1,1} C_{1,0,1}$$

Conditions [46] and [51] ([46] and [52]) are treated similarly using the corresponding boundary expressions. Thus,

$$[93] \quad c_h(X_0, 1, T) = c_h(X_0, 0, T) = \frac{1}{G_1} \left[\sum_{i=1}^{N+1} H_i c_h(X_i, R_0, T) + \sum_{i=1}^{N+1} C_{1,i} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) \right]$$

and

$$[94] \quad c_h(X_0, 1, T) = c_h(X_0, 0, T) = \frac{1}{G_1} \left[\sum_{i=1}^{N+1} H_i c_h(X_i, R_0, T) + \sum_{i=1}^{N+1} C_{1,i} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) \right]$$

Finally, the concentrations of species A at $R = 0$ and all X_i may be calculated by introducing [90] into [85]:

$$[95] \quad c_h(X_0, 0, T) = -\frac{1}{G_1} \left[\sum_{i=1}^{N+1} G_i c_h(X_i, R_0, T) + \sum_{i=1}^{N+1} C_{0,1,1} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) + C_{0,1,1} \sqrt{F} C_{0,1,1} \right]$$

while it holds from [93] and [94] ([94] and [97]), that

$$[96] \quad c_h(X_0, 0, T) = -\frac{1}{G_1} \left[\sum_{i=1}^{N+1} G_i c_h(X_i, R_0, T) + \sum_{i=1}^{N+1} C_{0,1,1} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) \right]$$

and that

$$[97] \quad c_h(X_0, 0, T) = -\frac{1}{G_1} \left[\sum_{i=1}^{N+1} G_i c_h(X_i, R_0, T) + \sum_{i=1}^{N+1} C_{0,1,1} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) \right]$$

Now we may substitute the unknown terms in eqs. [71]–[76]. From [71], using [90] and [95] it follows

$$[98] \quad \frac{dc_h}{dt} \Big|_{X_0} = \beta \left[R_{0,1,1} + \sum_{i=1}^{N+1} R_{1,i} c_h(X_i, R_0, T) \right] + \frac{1}{G_1} \left[I_{1,0} \sqrt{F} C_{0,1,1} + \sum_{i=1}^{N+1} I_{1,i} (\sqrt{F} C_{1,i} - C_{0,1,1} c_h(X_0, 1, T) - I_{1,0} C_{1,0,1} + I_{1,0} C_{1,0,1} c_h(X_0, 0, T)) + \sum_{i=1}^{N+1} I_{1,i} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) \right]$$

where

$$[99] \quad I_{1,0} = E_{0,1,1} C_{1,0} - E_{1,0} C_{0,1,1}$$

and

$$[100] \quad I_{1,i} = E_{1,i} C_{1,i} - E_{i,1} C_{1,i}$$

Combination of [72] with [80], [93], and [96] gives

$$[101] \quad \frac{dc_h}{dt} \Big|_{X_0} = \frac{\beta}{A_{1,1}} \left[\sum_{i=1}^{N+1} H_i c_h(X_i, R_0, T) - R_{0,1,1} \left(A_{1,0,1} + \sum_{i=1}^{N+1} A_{1,i} c_h(X_i, R_0, T) \right) + \frac{1}{G_1} \left[\sum_{i=1}^{N+1} I_{1,i} (\sqrt{F} C_{1,i} - C_{0,1,1} c_h(X_0, 1, T) - I_{1,0} C_{1,0,1} + I_{1,0} C_{1,0,1} c_h(X_0, 0, T)) + \sum_{i=1}^{N+1} I_{1,i} \sqrt{F} C_{1,i} c_h(X_i, R_0, T) \right] - \alpha c_h \right]$$

where

$$[102] \quad K_{i,j} = R_{i,j}A_{i,j} - R_{i,j}A_{i,j}$$

For the inner function describing the concentration of species C it holds from [73] with [81], [94] and [97]

$$[103] \quad \frac{dc_i}{dt} \Big|_{R_i} = \frac{\beta}{A_{i,j}} \left[\sum_{j=1}^{N_j} K_{i,j} c_i(X_i, R_i, T) \right] + \frac{1}{G_i} \left[\sum_{j=1}^{N_j} L_{i,j} \sqrt{D} C_{i,j} - C_{i,j+1} + L_{i,j} C_{i,j+1} + L_{i,j} C_{i,j+1} c_i(X_i, R_i, T) \right] + \sum_{j=1}^{N_j} L_{i,j} \sqrt{D} C_{i,j} c_i(X_i, R_i, T) + ac_i$$

The outer functions for all three species are derived following the same scheme. The time derivative of c_i in the outer domain is found from [74], [82], and [90], and in analogy the corresponding equations for species B or C may be calculated from [75], [83], and [93] or [76], [84], and [94].

$$[104] \quad \frac{dc_i}{dt} \Big|_{R_i} = \frac{\beta}{A_{i,j}} \left[K_{i,j+1} + \sum_{j=1}^{N_j} K_{i,j} c_i(X_i, R_i, T) \right] + \frac{\beta}{G_i} \left[L_{i,j+1} + \sum_{j=1}^{N_j} F_{i,j} K_{i,j} c_i(X_i, R_i, T) + \sum_{j=1}^{N_j} L_{i,j} c_i(X_i, R_i, T) \right] + ac_i$$

with

$$[105] \quad L_{i,j} = F_{i,j} G_i + C_{i,j} \sqrt{D} C_{i,j} F_{i,j}$$

$$[106] \quad \frac{dc_i}{dt} \Big|_{R_i} = \frac{\beta}{A_{i,j}} \left[\sum_{j=1}^{N_j} K_{i,j} c_i(X_i, R_i, T) \right] + \frac{\beta}{G_i} \left[\sum_{j=1}^{N_j} F_{i,j} K_{i,j} c_i(X_i, R_i, T) + \sum_{j=1}^{N_j} L_{i,j} c_i(X_i, R_i, T) \right] + ac_i$$

$$[107] \quad \frac{dc_i}{dt} \Big|_{R_i} = \frac{\beta}{A_{i,j}} \left[\sum_{j=1}^{N_j} K_{i,j} c_i(X_i, R_i, T) \right] + \frac{\beta}{G_i} \left[\sum_{j=1}^{N_j} F_{i,j} K_{i,j} c_i(X_i, R_i, T) + \sum_{j=1}^{N_j} L_{i,j} c_i(X_i, R_i, T) \right] + ac_i$$

Now the set of six second order partial differential equations [26]–[28] and [30]–[32] has been transformed into a set of $3N_j(N_j + N_s)$ simultaneous ordinary differential equations [90], [101], [103], [104], [106], and [107]. Such systems may be easily integrated by computer programs to give the time dependence of the concentrations of species A, B, and C at the collocation points (X_i, R_i) and (X_i, R_i) . We describe the concentration profiles which develop in front of the electrode by the concentrations at the collocation points.

We note the structural similarity of the formulae describing the time derivatives of the concentrations. This is one of the advantages of the orthog-

onal collocation method, for if the theory has been derived once it is very easy to express it in a computer language. Furthermore, if more complicated mechanisms are to be simulated, only minor changes in the formulae have to be performed. Also, the extension of the theory to other experimental techniques, i.e. other boundary conditions, is easy. This will be shown in another paper of this series.

We also note that the implementation of the expressions describing species C is not essential to calculate the concentration profiles of A and B, for C is not electroactive and does not react to or with an electroactive intermediate [16]. However, if one

desires, for example, to calculate the spectroelectrochemical response [14] of species C one is interested in the concentration profiles of C as well.

Calculation of the current

In the one-dimensional diffusion model the current i is simply calculated from the flux of the substrate at the electrode surface:

$$[108] \quad i = nFAD \left(\frac{\partial c_A}{\partial x} \right)_{x=0}$$

where A is the electroactive area of the electrode. This implies that the flux of the substrate is uniform over A .

This is clearly not true in the case of a two-dimensional model considering edge effects. Instead, the current is given by

$$[109] \quad i = nFD \int_0^{2\pi} \left(\frac{\partial c_A}{\partial r} \right)_{r=0} dr$$

where the r -dependent flux through an infinitesimally small part $2\pi r dr$ of the electrode is integrated over the electroactive section, $0 \leq r \leq r_0$ of the r -coordinate.

Using the transformation equations [21]–[23], [25], and the definition of β (eq. [29]), eq. [109] yields

$$[110] \quad i = \frac{nFD \int_0^{2\pi} \left(\frac{\partial c_A}{\partial x} \right)_{x=0} dR}{L} = 2nFAC \frac{\sqrt{D}}{\sqrt{t}} \int_0^1 R \left(\frac{\partial c_A}{\partial X} \right)_{X=0} dR$$

The chronoamperometric "constant" $i\sqrt{t}$ is given by

$$[111] \quad i\sqrt{t} = 2nFAC \frac{\sqrt{D}}{\sqrt{t}} \int_0^1 R \left(\frac{\partial c_A}{\partial X} \right)_{X=0} dR$$

Alternatively, we may express this product in terms of the "Cottrell constant" (2)

$$[112] \quad (i\sqrt{t})_C = \frac{nFAC \sqrt{D}}{\sqrt{t}}$$

as

$$[113] \quad \frac{i\sqrt{t}}{(i\sqrt{t})_C} = 2\sqrt{\pi} \int_0^1 R \left(\frac{\partial c_A}{\partial X} \right)_{X=0} dR$$

Since the differential quotients $(\partial c_A / \partial X)$ are given at the collocation points we may calculate the integral exactly by a quadrature formula of the type (6), [20], p. 134)

$$[114] \quad \int_0^1 R \left(\frac{\partial c_A}{\partial X} \right)_{X=0} dR = \sum_{j=1}^{N_j} Q_j R_j$$

where the Q_j are elements of a vector depending on the trial function chosen.

Thus, from [67]

$$[115] \quad \int_0^1 R \left(\frac{\partial c_A}{\partial X} \right)_{X=0} dR = \sum_{j=1}^{N_j} Q_j \left[R_j \sum_{i=1}^{N_j} A_{i,j} c_i(X_i, R_i, T) \right]$$

Expanding the sum over j and introducing the boundary conditions [36] and [42b], [115] changes to

$$[116] \quad \int_0^1 R \left(\frac{\partial c_A}{\partial X} \right)_{X=0} dR = \sum_{j=1}^{N_j} Q_j \left[R_j A_{i,j+1} + R_j \sum_{i=1}^{N_j} A_{i,j} c_i(X_i, R_i, T) \right]$$

The Gaussian weights Q_j for $j=1$ and $j=N_j+2$ are equal to zero, thus [116] reduces to

$$[117] \quad \int_0^1 R \left(\frac{\partial c_A}{\partial X} \right)_{X=0} dR = \sum_{j=1}^{N_j} Q_j \left[R_j A_{i,j+1} + R_j \sum_{i=1}^{N_j} A_{i,j} c_i(X_i, R_i, T) \right]$$

which is a convenient form for computation of the integral.

Acknowledgment

We thank the Office of Naval Research, Washington, D.C., for support of part of this work.

1. P. J. LINGANE, *Anal. Chem.* **30**, 1733 (1958).
2. D. D. McDONALD, *Transient techniques in electrochemistry*, Plenum Press, New York and London, 1977, p. 72.
3. R. LÖNN and V. D. PAMAR, *Acta Chem. Scand.* **31**, 349 (1977).
4. R. S. MCDONALD and I. SMITH, *Anal. Chem.* **30**, 765 (1958).
5. Z. G. SONG and P. J. LINGANE, *J. Phys. Chem.* **68**, 4821 (1964).
6. J. B. FLANNAGAN and L. MARCOU, *J. Phys. Chem.* **77**, 1051 (1973).
7. B. SPEISER and A. RABIN, *Electrochim. Acta*, **23**, 953 (1978).
8. J. HAYES, *Rev. Roum. Phys. Chem.* **34**, 783 (1989).
9. J. HAYES, *J. Electroanal. Chem.* **126**, 75 (1981).

10. K. B. OGDON, *J. Electroanal. Chem.* **122**, 1 (1981).
11. K. AOKI and J. OHTSUKA, *J. Electroanal. Chem.* **122**, 19 (1981).
12. L. F. WATKINS and P. W. CARR, *J. Electroanal. Chem.* **81**, 1 (1977).
13. A. BEWICK, J. MOLLAN, and S. PONS, *Electrochim. Acta*, **25**, 77 (1979).
14. A. BEWICK, J. MOLLAN, and S. PONS, *Electrochim. Acta*, **28**, 911 (1983).
15. S. PONS and P. P. SCHMIDT, *Electrochim. Acta*, **28**, 927 (1983).
16. B. SUNDEN and A. RIGG, *J. Electroanal. Chem.* **105**, 1 (1979).
17. B. SUNDEN, *J. Electroanal. Chem.* **110**, 69 (1980).
18. S. PONS, *Can. J. Chem.* **59**, 1518 (1981).
19. B. A. FINLAYSON, in *Mathematics in science and engineering*, Vol. 87, Edited by R. Bellman, Academic Press, New York and London, 1972, p. 107.
20. J. VILKADIN and M. U. LILJA, 1981, *Solutions of differential equations models by polynomial approximation*, Prentice Hall, Englewood Cliffs, 1978.

Correlation of the hydrogenation catalytic activity of dinuclear complexes of rhodium(I) with the formation of carbonyl-bridged complexes

ALAN R. SANDER

Alberta Research Council, 11113-87 Avenue, Edmonton, Alta., Canada T6G 2C7

Received May 8, 1981

ALAN R. SANDER, *Can. J. Chem.* **60**, 1363 (1982).

Methanolic solutions of the dinuclear cationic complexes of rhodium(I), $[\text{Rh}_2(\mu\text{-X})(\text{CO})_2(\text{Ph})_2\text{ECN}(\text{EPPh}_2)]^+$ ($\text{X} = \text{Cl}$, Br; $\text{E} = \text{P}$, As) and iridium(I), $[\text{Ir}_2(\text{Cl})_2(\text{Ph})_2\text{PCH}_2\text{PPh}_2]_2^{2+}$, are each active catalysts for the hydrogenation of alkynes to alkenes, and alkenes to alkanes. Neutral complexes with pseudohalogen ligands have also been investigated but only the cyano-complex $[\text{Rh}_2(\mu\text{-N})_2(\text{CO})_2(\text{Ph})_2\text{PCH}_2\text{PPh}_2]_2$, and the arsenic analog, are of significant catalytic activity. Only those complexes which reversibly form bridged complexes with ligands such as CO are also catalytically active.

Similar complexes with dihalogen ligands are not efficient catalysts under the conditions tested.

ALAN R. SANDER, *Can. J. Chem.* **60**, 1363 (1982).

Les complexes cationiques dinucléaires de rhodium(I), $[\text{Rh}_2(\mu\text{-X})(\text{CO})_2(\text{Ph})_2\text{ECN}(\text{EPPh}_2)]^+$ ($\text{X} = \text{Cl}$, Br; $\text{E} = \text{P}$, As), et d'iridium(I), $[\text{Ir}_2(\text{Cl})_2(\text{Ph})_2\text{PCH}_2\text{PPh}_2]_2^{2+}$, en solution dans le méthanol, sont des catalyseurs actifs d'hydrogénation des alcynes en alcènes et d'alcènes en alcanes. On a également étudié les complexes neutres ayant des ligands pseudohalogénés, mais seuls les complexes de type cyano $[\text{Rh}_2(\mu\text{-N})_2(\text{CO})_2(\text{Ph})_2\text{PCH}_2\text{PPh}_2]_2$ et leurs analogues de type arsène ont une activité catalytique significative. Seuls les complexes qui forment de manière réversible des complexes liés par un pont avec des ligands, tels le CO, sont également des catalyseurs actifs.

Les complexes similaires avec les ligands dihalogénés ne sont pas des catalyseurs efficaces dans les conditions étudiées.

(Traduit par le journal)

Introduction


The first dinuclear rhodium complex of the geometry designated "A-frame" (1) $[\text{Rh}_2(\mu\text{-S})(\text{CO})_2(\text{dppm})_2]$ (dppm is $\text{Ph}_2\text{PCH}_2\text{PPh}_2$ and dpm is $\text{Ph}_2\text{AsCH}_2\text{AsPh}_2$) was described in 1977 (1). The synthesis and structure of related complexes of rhodium or iridium has since received much attention (2-16). For complexes of rhodium(I), several instances of co-ordination of two-electron donor ligands (L; e.g. CO, SO_2) at the site bridging the metal centers (2) have been reported (2, 3, 6, 9, 12-16). The complex cation, $[\text{Rh}_2(\mu\text{-Cl})(\text{CO})_2(\text{dppm})_2]^+$, forms a 1:1 labile, symmetrical (μ , ^{31}P NMR) complex with diphenylacetylene (6),¹ and similar neutral complexes have also recently been described (11). Co-ordination of an alkyne at the bridging site (2; L = alkyne) is accompanied by lengthening of the carbon-carbon bond such that the product is accurately described as aryl-dimetallated alkynyl complex (11). No complex in which an alkene is co-ordinated at either a bridging or terminal site of an A-frame complex has yet been described.

¹Revised received February 12, 1982.

²This system has been investigated elsewhere (16) Canada, private communication) and in this laboratory. The differences in spectroscopic parameters between solutions of the cationic and neutral complexes are now considered to be due to differences in conditions. A cationic complex containing bridging $\text{CH}_3\text{XCOXCH}_3$ has been well characterized (16).



1 ($\text{E} = \text{P}$ or As, X is no longer ligand of formal charge +). Y is a second two-electron donor ligand, usually CO



The A-frame structure, with the proximity of the two metal centres, has strong potential for dinuclear activation of unsaturated substrates. However, the complex $[\text{Rh}_2(\mu\text{-S})(\text{CO})_2(\text{dppm})_2]$ has been shown to have low activity for the hydrogenation of ethylene (13). Herein I report that those cationic or neutral complexes of rhodium (or iridium) which form carbonyl-bridged complexes (2, L = CO) are also efficient catalysts.

TECHNICAL REPORT DISTRIBUTION LIST, 359

No. Copies	No. Copies
Dr. R. Mousk Naval Research Laboratory Code 6130 Washington, D.C. 20375	1
Dr. John F. Howlhan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1
Dr. D. P. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1
Dr. Alan Bewick Department of Chemistry The University Southampton, 509 3NW England	1
Dr. A. Niy NAVSEA-5433 WC #4 2341 Jefferson Davis Highway Arlington, Virginia 20362	1
Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, DC 20376	1
M. L. Robertson Manager, Electrochemical Power Sources Division Naval Weapons Support Center Crane, Indiana 47522	1
Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720	1
Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709	1
Dr. Denton Elliott Air Force Office of Scientific Research Building APB Washington, DC 20332	1
Dr. David Aikens Chemistry Department Rensselaer Polytechnic Institute Troy, NY 12181	1
Dr. A. F. S. Lever Chemistry Department York University Downsview, Ontario M3J1P3 Canada	1
Mr. Maurice F. Murphy Naval Sea Systems Command 61832 2221 Jefferson Davis Highway Arlington, VA 20360	1
Dr. Stanislaw Szpak Naval Ocean System Center Code 6343 San Diego, CA 95152	1
Dr. Gregory Farrington Department of Materials Science & Engineering University of Pennsylvania Philadelphia, PA 19104	1
Dr. Bruce Dunn Department of Engineering & Applied Science University of California Los Angeles, CA 90024	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

No. Copies	No. Copies
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1
Larry E. Flew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1
S. Rubw DOE (STOR) 600 E Street Washington, D.C. 20545	1
Dr. Aaron Mold Brown University Department of Chemistry Providence, Rhode Island 02192	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1
Dr. N. W. Nicholson Electronics Research Center Rockwell International 3370 Wilshire Avenue Anaheim, California	1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. R. P. Van Duren Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. E. Stanley Fenn Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6C 2G7	1
Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
Dr. P. David Bach EIC Corporation 55 Canal Street Newton, Massachusetts 02158	1
Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 3NW England	1
Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Mr. James R. Haden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1

SP472-3/A1

472:GAN:716:enJ
78u472-608

SP472-3/A11

472:GAN:716:64c
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies	No. Copies
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRO-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709 1
ONR Western Regional Office Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152 1
ONR Eastern Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 606 Summer Street Boston, Massachusetts 02210	1	Naval Weapons Center Attn: Dr. A. S. Amster, Chemistry Division China Lake, California 93555 1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401 1
The Assistant Secretary of the Navy (REAS) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940 1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380 1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22304	12	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401 1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 92132 1
		Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112 1

TECHNICAL REPORT DISTRIBUTION LIST, 350

	No. Copies	No. Copies
Dr. Paul Delaney Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Headra Department of Chemistry University of Southampton Southampton SO9 5NR United Kingdom 1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 44106	1	Dr. Sam Perone Department of Chemistry Purdue University West Lafayette, Indiana 47907 1
Dr. D. M. Bannion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce M. Murver Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514 1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152 1
Dr. J. J. Auborn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Warfare Center White Oak Laboratory Silver Spring, Maryland 20910 1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201 1
Dr. T. Katan Lockheed Missiles & Space Co., Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochemical Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040 1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063 1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1	Dr. H. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181 1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Arlington, Massachusetts 01803	1	

SP472-3/B17

472:GAN:716:1ab
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, 359

No.
Copies

Dr. Micha Tomkiewicz
Department of Physics
Brooklyn College
Brooklyn, NY 11210

1

Dr. Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras, PR 00931

1

Dr. Joseph Gordon II
IBM Corporation
K33/281
5600 Cottle Road
San Jose, CA 95193

1

Dr. Robert Somoano
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, CA 91103

1

END

DATE
FILMED

9 82

DTIC